

11. Behandling ved ukendt primær tumor

Tina Tos, Krzysztof T. Drzewiecki

Ansvarlig: Tina Tos. Opdateret: 13-02-2012

Behandling ved ukendt primær tumor følger den anbefalede behandling for locoregional recidiv samt metastasering med kendt primær melanom. Nogle studier har vist en lidt bedre overlevelse hos patienter med ukendt primær tumor.[1-4]

Reference List

1. Cormier JN, Xing Y, Feng L, Huang X, Davidson L, Gershenwald JE *et al.* Metastatic melanoma to lymph nodes in patients with unknown primary sites. *Cancer* 2006; **106**(9): 2012-2020.
2. Lee CC, Faries MB, Wanek LA, Morton DL. Improved survival after lymphadenectomy for nodal metastasis from an unknown primary melanoma. *J Clin Oncol* 2008; **26**(4): 535-541.
3. Lee CC, Faries MB, Wanek LA, Morton DL. Improved survival for stage IV melanoma from an unknown primary site. *J Clin Oncol* 2009; **27**(21): 3489-3495.
4. Vijuk G, Coates AS. Survival of patients with visceral metastatic melanoma from an occult primary lesion: a retrospective matched cohort study. *Ann Oncol* 1998; **9**(4): 419-422.